

The Relationship Between Noise Correlation and the Green's Function in the Presence of Degeneracy and the Absence of Equipartition

Victor C. Tsai¹

¹ *Geologic Hazards Science Center, United States Geological Survey, Golden, CO 80401, USA.
Email: vtsai@post.harvard.edu*

Received 2010 June 4; in original form 2010 February 23

SUMMARY

Recent derivations have shown that when noise in a physical system has its energy equipartitioned into the modes of the system, there is a convenient relationship between the cross correlation of time series recorded at two points and the Green's function of the system. Here, we show that even when energy is not fully equipartitioned and modes are allowed to be degenerate, a similar (though less general) property holds for equations with wave equation structure. This property can be used to understand why certain seismic noise correlation measurements are successful despite known degeneracy and lack of equipartition on the Earth.

Key words: Theoretical seismology; statistical seismology; wave propagation; surface waves and free oscillations; seismic tomography

1 INTRODUCTION

Beginning with the work of Lobkis & Weaver (2001), a number of authors have shown that under very special circumstances, there exists a convenient relationship between the cross correlation of time series recorded at two points and the Green's function of the system. The assumptions made in the various derivations ranges substantially, with some authors assuming that the energy in normal modes is uncorrelated and perfectly equipartitioned (Lobkis & Weaver 2001), some assuming an isotropic energy flux (Snieder 2004; Sanchez-Sesma et al. 2006; Sanchez-Sesma & Campillo 2006; Tsai 2009), and others relying on reciprocity relations (Wapenaar 2004; Wapenaar et al. 2006).

In order to make use of these properties, it is further often assumed that certain systems, like the Earth, satisfy all of the assumptions of the theories. However, some assumptions are clearly not satisfied, bringing into question the applicability of the theories. For example, it is well known that most ambient seismic noise is generated at the surface and that modes with higher sensitivity near the surface will be preferentially excited, resulting in a lack of equipartition; it is also known that microseisms are primarily generated in certain oceanic regions and therefore do not provide an isotropic incidence of energy flux. Do relationships between the cross correlation and Green's function still exist under less restrictive conditions? The relationship between the assumptions is also not entirely clear. For example, are the assumptions of equipartition and that of isotropic energy flux equivalent? In this work, we

address some of these questions. Using a normal mode approach, we find important differences in the cross correlation-Green's function relationship when modes are equipartitioned or not and when modes are degenerate or not. We then show a few simple examples of applying these relationships.

2 DEFINITIONS

The approach taken in this work is to consider a mathematical system in which wave-like modes exist [as in Lobkis & Weaver (2001)] and determine the consequences of the system satisfying various assumptions. Specifically, we assume that the system of interest can be written in the form

$$\frac{\partial^2 u(\mathbf{x}, t)}{\partial t^2} = \mathcal{L}[u(\mathbf{x}, t)], \quad (1)$$

where \mathbf{x} is a spatial variable, t is time, and \mathcal{L} is a spatial operator with given boundary conditions such that there is a basis of spatial eigenfunctions (normal modes), $s_k(\mathbf{x})$. That is, we assume that

$$\mathcal{L}[s_k(\mathbf{x})] = \lambda_k s_k(\mathbf{x}) = -\omega_k^2 s_k(\mathbf{x}), \quad (2)$$

where the eigenfrequencies ω_k are real, $s_k(\mathbf{x})$ are orthonormal, satisfying $\langle s_k, s_{k'} \rangle = \delta_{kk'}$, and any arbitrary function $f(\mathbf{x})$ in the solution space can be expressed as

$$f(\mathbf{x}) = \sum_k A_k s_k(\mathbf{x}). \quad (3)$$

In the above, $\langle \cdot, \cdot \rangle$ denotes the appropriate inner product, δ_{ij} is the Kronecker delta, and the sum on k may (or may not) be infinite. One should note that the acoustic wave equation and elastic wave equation satisfy these assumptions, regardless of the number of spatial dimensions. One may also note that these 2 equations do not include attenuation.

Considering the previous assumptions, standard separation of variables then yields the general solution

$$u(\mathbf{x}, t) = \sum_k a_k s_k(\mathbf{x}) \cos(\omega_k t + \phi_k), \quad (4)$$

where a_k and ϕ_k are constants determined by the initial conditions. We define two modes l and m to be equipartitioned if $a_l = a_m$.

The Green's function, $G(\mathbf{x}, t; \mathbf{x}_0, t_0)$, for an impulse at location \mathbf{x}_0 and time t_0 , satisfies

$$\frac{\partial^2 G}{\partial t^2} - \mathcal{L}[G] = \delta(\mathbf{x} - \mathbf{x}_0) \delta(t - t_0), \quad (5)$$

where $\delta(x)$ is the Dirac delta function. Choosing $t_0 = 0$ for convenience, G can be written as a modal sum as in Eq. (4), with coefficients $a_k = s_k(\mathbf{x}_0)/\omega_k$, $\phi_k = -\pi/2$ when $t > 0$. Writing this out explicitly yields

$$G(\mathbf{x}, t; \mathbf{x}_0, 0) = \begin{cases} \sum_k \frac{1}{\omega_k} s_k(\mathbf{x}) s_k(\mathbf{x}_0) \sin \omega_k t & \text{if } t \geq 0 \\ 0 & \text{if } t < 0 \end{cases}. \quad (6)$$

We further define the extended Green's function as

$$\begin{aligned} G^{Ex}(\mathbf{x}, t; \mathbf{x}_0, 0) &\equiv G(\mathbf{x}, t; \mathbf{x}_0, 0) - G(\mathbf{x}, -t; \mathbf{x}_0, 0) \\ &= \sum_k \frac{1}{\omega_k} s_k(\mathbf{x}) s_k(\mathbf{x}_0) \sin \omega_k t, \end{aligned} \quad (7)$$

for all t . The time derivative is then given by

$$\frac{dG^{Ex}}{dt} = \sum_k s_k(\mathbf{x}) s_k(\mathbf{x}_0) \cos \omega_k t. \quad (8)$$

Finally, we define the normalized cross correlation as

$$C_{fg}^T(t) \equiv \frac{1}{2T} \int_{-T}^T f(\tau) g(\tau + t) d\tau, \quad (9)$$

and the limiting value as

$$C_{fg}(t) \equiv f \star g(t) \equiv \lim_{T \rightarrow \infty} C_{fg}^T(t). \quad (10)$$

For this choice, cross correlations between (infinitely long) sinusoidal signals are well defined. An important example of this, which is used throughout this work, is when $f(t) = \cos(\omega_k t + \phi_k)$ and $g(t) = \cos(\omega_{k'} t + \phi_{k'})$. For this choice of f and g , direct calculation yields

$$C_{fg}(t) = \begin{cases} \frac{1}{2} \cos(\omega_k t + \phi_{k'} - \phi_k) & \text{if } \omega_k = \omega_{k'} \\ 0 & \text{if } \omega_k \neq \omega_{k'} \end{cases}. \quad (11)$$

In the following sections, the subscript ' fg ' on cross correlations will refer to the locations of the recorded time series (rather than functions).

3 RELATIONSHIPS BETWEEN THE CROSS CORRELATION AND THE GREEN'S FUNCTION

We now provide the relationship between cross correlations and the Green's function under various assumptions.

3.1 Equipartitioned, Non-Degenerate Modes

In this most restricted case, it is assumed that energy is equipartitioned between all modes so that all modes have equal modal amplitudes, $a_k = \text{const.} = A$, so that $u(\mathbf{x}, t)$ can be written as

$$u(\mathbf{x}, t) = A \sum_k s_k(\mathbf{x}) \cos(\omega_k t + \phi_k). \quad (12)$$

It is also assumed that no modes are degenerate, i.e. $\omega_k \neq \omega_{k'}$ when $k \neq k'$. With these assumptions, applying Eq. (11) term by term yields

$$C_{x_1 x_2}(t) = \frac{A^2}{2} \sum_k s_k(\mathbf{x}_1) s_k(\mathbf{x}_2) \cos \omega_k t. \quad (13)$$

Comparing Eq. (13) with Eq. (8), one immediately finds that

$$C_{x_1 x_2}(t) = \frac{A^2}{2} \frac{dG^{Ex}(\mathbf{x}_1, t; \mathbf{x}_2, 0)}{dt}, \quad (14)$$

i.e. the cross correlation is identical to the time derivative of the Green's function, up to an amplitude factor. Expressions similar to Eq. (14) have been previously noted by a number of authors including Lobkis & Weaver (2001), Wapenaar (2004) and Roux et al. (2005), though not with the same assumptions. Importantly, this relationship holds for a completely deterministic process in which the phases, ϕ_k , are arbitrary but unchanging and no assumption regarding the randomness of these phases through time is required.

3.2 Equipartitioned, Degenerate Modes

Degeneracy of modes, with $\omega_k = \omega_{k'}$ for some $k \neq k'$, is common in many systems. For example, symmetric or periodic boundary conditions often leads to degeneracy. When modes are allowed to be degenerate, Eq. (13) no longer holds because of the existence of cross terms between degenerate modes. In order for these terms to disappear, one must make the additional assumption (as in Lobkis & Weaver (2001)) that there is an ensemble of states in which the phases, ϕ_k , are randomly chosen in each realization or, equivalently, that the ϕ_k are periodically reset to random values. To demonstrate the result of this additional assumption, we take the ensemble point of view, assuming there exist N different realizations of Eq. (12) with randomly chosen ϕ_k so that

$$u_j(\mathbf{x}, t) = A \sum_k s_k(\mathbf{x}) \cos(\omega_k t + \phi_{jk}), \quad (15)$$

where ϕ_{jk} are randomly chosen, $1 \leq j \leq N$. The ensemble cross correlation C_{fg}^E is set equal to the average of the cross correlations for these N realizations. Assuming that modes l and m are degenerate ($\omega_l = \omega_m$), then the part of $C_{x_1 x_2}^E$ due to just those terms would be

$$\begin{aligned} C_{x_1 x_2}^{Elm} &= \frac{A^2}{2} \left[s_l(\mathbf{x}_1) s_l(\mathbf{x}_2) \cos(\omega_l t) + s_m(\mathbf{x}_1) s_m(\mathbf{x}_2) \cos(\omega_m t) \right. \\ &\quad + \frac{1}{N} s_l(\mathbf{x}_1) s_m(\mathbf{x}_2) \cos(\omega_l t + \phi_{1m} - \phi_{1l}) \\ &\quad \left. + \frac{1}{N} s_l(\mathbf{x}_1) s_m(\mathbf{x}_2) \cos(\omega_l t + \phi_{2m} - \phi_{2l}) + \dots \right]. \end{aligned} \quad (16)$$

If $\phi_{jm} - \phi_{jl}$ are randomly distributed (i.e. uncorrelated), then the sum of the cross-terms will grow as \sqrt{N} , so that Eq. (16) reduces

to

$$C_{x_1 x_2}^{Elm} = \frac{A^2}{2} \left[s_l(\mathbf{x}_1) s_l(\mathbf{x}_2) \cos(\omega_l t) + s_m(\mathbf{x}_1) s_m(\mathbf{x}_2) \cos(\omega_m t) \right. \\ \left. + \frac{1}{\sqrt{N}} s_l(\mathbf{x}_1) s_m(\mathbf{x}_2) \cos(\omega_l t + \phi_{lm}^{avg}) + \dots \right], \quad (17)$$

where ϕ_{lm}^{avg} is the average of $\phi_{jm} - \phi_{jl}$ (which can be calculated by summing phases in the complex plane). As $N \rightarrow \infty$, then again the cross terms become vanishingly small

$$\lim_{N \rightarrow \infty} C_{x_1 x_2}^{Elm} = \frac{A^2}{2} \left[s_l(\mathbf{x}_1) s_l(\mathbf{x}_2) \cos(\omega_l t) \right. \\ \left. + s_m(\mathbf{x}_1) s_m(\mathbf{x}_2) \cos(\omega_m t) \right]. \quad (18)$$

Since the same argument can be made for all pairs of degenerate modes, we retrieve the relationship

$$\lim_{N \rightarrow \infty} C_{x_1 x_2}^E(t) = \frac{A^2}{2} \frac{dG^{Ex}(\mathbf{x}_1, t; \mathbf{x}_2, 0)}{dt}. \quad (19)$$

Thus, we observe that only when ϕ_{jk} sample sufficiently many states does the cross correlation converge upon the Green's function. This result is similar to that of Lobkis & Weaver (2001).

For a system with attenuation, the ϕ_k [e.g. of Eq. (12)] are naturally reset to random values after an e-folding attenuation time as long as there exists random forcing through time (since the effects of forcing from times sufficiently far in the past are attenuated away). This is an example of a case in which the periodic resetting of phases results in an effective sampling of different ensemble states. In such a case, the number of realizations, N , would roughly be equivalent to the number of e-folding attenuation times in the time series used. Provided that Eq. (11) is well approximated within the e-folding time (i.e. attenuation is weak), then we again achieve Eq. (19) as a result.

3.3 Non-Equipartitioned, Non-Degenerate Modes

We now relax the assumption of equipartition but consider the case where there are no degenerate modes. Again applying Eq. (11) term by term yields

$$C_{x_1 x_2}(t) = \sum_k \frac{a_k^2}{2} s_k(\mathbf{x}_1) s_k(\mathbf{x}_2) \cos \omega_k t. \quad (20)$$

Comparing Eq. (20) with Eq. (8), we observe that while the summed expressions cannot be compared directly, each term is still equivalent up to an amplitude factor, $a_k^2/2$, which can be different for each k . In many applications, including the seismological one, 'phase' or 'travel-time' information is useful regardless of whether amplitude measurements can be made easily (whether due to instrumental limitations or structural complexities). (It is perhaps useful to note that if one only has band-limited data, the full Green's function can never be reconstructed anyway.) Thus, we observe that the cross correlation still yields a useful portion of the Green's function. Specifically, if one filters both the cross correlation and the Green's function at frequency ω_k , then one is left with only the k th term of the sum, hence

$$\mathcal{F}_{\omega_k}[C_{x_1 x_2}(t)] = A_k \mathcal{F}_{\omega_k} \left[\frac{dG^{Ex}(\mathbf{x}_1, t; \mathbf{x}_2, 0)}{dt} \right], \quad (21)$$

where \mathcal{F}_{ω} is a perfect filter at frequency ω and $A_k = a_k^2/2$. A similar relationship was previously drawn by Sanchez-Sesma &

Campillo (2006) for the less general case of propagating P, SV and SH waves.

3.4 Non-Equipartitioned, Degenerate Modes

Finally, we relax both the assumption of equipartition and of non-degeneracy. Following the procedure of Section 3.2, we can still apply Eq. (11) and obtain

$$\lim_{N \rightarrow \infty} C_{x_1 x_2}^E(t) = \sum_k \frac{a_k^2}{2} s_k(\mathbf{x}_1) s_k(\mathbf{x}_2) \cos \omega_k t. \quad (22)$$

As in Section 3.3, there is term-by-term equality of the cross correlation and the Green's function. However, if degenerate modes have different relative amplitudes (i.e. $a_l \neq a_m$ when $\omega_l = \omega_m$), then even a filtered response will not yield results equivalent up to an amplitude scale factor.

This analysis shows that the necessary condition on (perfect) retrieval of Green's function 'phase' information using cross correlations is that all degenerate modes are equipartitioned. The lack of equipartition between non-degenerate modes only results in a loss of amplitude information. When this equipartition of degenerate modes exists, then

$$\mathcal{F}_{\omega_k} \left[\lim_{N \rightarrow \infty} C_{x_1 x_2}^E(t) \right] = A_k \mathcal{F}_{\omega_k} \left[\frac{dG^{Ex}(\mathbf{x}_1, t; \mathbf{x}_2, 0)}{dt} \right], \quad (23)$$

where the $A_k = a_k^2/2$ are the same for each degeneracy group.

It may further be observed that if a subset of these degenerate modes, s_l (with $l \in S$), are equipartitioned with amplitude a_l , then it may be possible to partially retrieve Green's function 'phase' information. This would be possible, for example, if the arrival phases of all other degenerate modes s_m ($m \notin S$) are cleanly separated in time from modes s_l such that a time windowing of both sides of Eq. (23) results in picking out only modes s_l . Since windowing introduces neighboring frequencies, this partial retrieval would inherently make use of approximately degenerate modes (and their group arrivals). This case of partially equipartitioned modes may apply in many cases. For example, on the Earth, a subset of fundamental-mode Rayleigh waves may be approximately equipartitioned while all other modes (including higher-order modes) may not. On the Earth, higher-order modes are typically poorly excited (most forcing is near-surface and excites fundamental-mode waves more easily), resulting in much lower modal amplitudes (a_k) compared to the fundamental-mode amplitudes. This can therefore help explain why phase information for fundamental-mode Rayleigh waves is typically easily obtained, but similar phase information for higher-order (body wave) modes is difficult to obtain.

4 EXAMPLES

In this section, three examples are given of systems in which the results of Section 3 can be applied. In the first example, the simple case of waves on a periodic string, many of the properties discussed in Section 3 can be easily understood. Prior to discussing the third example of waves on the Earth, we provide a second example of waves on a periodic square. This example has many similarities to the Earth case and has results that can be generalized to the Earth, yet is simpler to understand.

4.1 Waves on a Periodic String of Length L

A string of length L with periodic boundary conditions is an example of a simple system in which waves have degeneracy. In this case, $\mathcal{L} = c^2 \partial^2 / \partial x^2$, and for each $\omega_k = k\pi c/L$ (where k is an integer, as before) there are two eigenfunctions $s_{k1}(x) = \sin(k\pi x/L)$ and $s_{k2}(x) = \cos(k\pi x/L)$ for $1 \leq k < \infty$, each with 2 initial conditions. By the results of Section 3.4, we know that if each pair of degenerate modes has equal amplitudes, then Eq. (23) holds.

Equivalently, one can express these 4 (standing wave) modes as 4 traveling waves, two of which travel in the positive x direction, two of which travel in the negative direction. Written as traveling waves

$$\begin{aligned} & a_{k1} \sin(\kappa x) \cos(\omega t + \phi_{k1}) + a_{k2} \cos(\kappa x) \cos(\omega t + \phi_{k2}) \\ = & \left[-\frac{a_{k1}}{2} \sin(\omega t + \phi_{k1} - \kappa x) + \frac{a_{k2}}{2} \cos(\omega t + \phi_{k2} - \kappa x) \right] \\ & + \left[\frac{a_{k1}}{2} \sin(\omega t + \phi_{k1} + \kappa x) + \frac{a_{k2}}{2} \cos(\omega t + \phi_{k2} + \kappa x) \right]. \quad (24) \end{aligned}$$

If, now, one assumes that $-a_{k1} + a_{k2}$ and $a_{k1} + a_{k2}$ are equal averaged over an ensemble (as in Section 3.2, corresponding to equal contributions from waves propagating in the positive and negative directions), then

$$\begin{aligned} & \lim_{N \rightarrow \infty} C_{x_1 x_2}^{E k_1 k_2} \\ = & \frac{A}{2} \left[\cos[\omega t + \kappa(x_1 - x_2)] + \cos[\omega t - \kappa(x_1 - x_2)] \right] \\ = & A \left[\cos(\kappa x_1) \cos(\kappa x_2) + \sin(\kappa x_1) \sin(\kappa x_2) \right] \cos(\omega t). \quad (25) \end{aligned}$$

We therefore again observe that Eq. (23) holds. In other words, having an ensemble of traveling waves that on average has equal amplitudes is equivalent to having an ensemble of standing-wave modes with equal amplitudes. In this case, having equal amplitude traveling waves corresponds to an isotropic density of sources (i.e. an equal amount of incident wave energy arriving from positive and negative x). Thus, there is an equivalence between equipartition amongst degenerate modes and an ‘isotropic’ incidence of waves. One should note that non-degenerate modes need not be equipartitioned in order for ‘isotropy’ to exist.

On the other hand, even if the negative and positive arriving waves are not equipartitioned, then

$$\begin{aligned} & \lim_{N \rightarrow \infty} C_{x_1 x_2}^{E k_1 k_2} = \\ & A \cos[\omega t + \kappa(x_1 - x_2)] + B \cos[\omega t - \kappa(x_1 - x_2)]. \quad (26) \end{aligned}$$

If one can pick out only the positive arrival (e.g. by time windowing the signal, as described in Section 3.4) then one can still identify the phase $\omega t - \kappa(x_1 - x_2)$ and hence determine a phase travel time ω/κ .

4.2 Waves on a Periodic Square of Size L

In this case, $\mathcal{L} = c^2(\partial^2/\partial x^2 + \partial^2/\partial y^2)$ and the eigenfrequencies can be expressed as $\omega_{lm}^2 = \pi^2 c^2(l^2 + m^2)/L^2$ for integer l and m . Periodic boundary conditions imply that each ω_{lm} has 4 modes given by all products of $\{\sin(\kappa_l x), \cos(\kappa_l x)\}$ and $\{\sin(\kappa_m y), \cos(\kappa_m y)\}$. Furthermore, there is additional degeneracy due to the fact that $l^2 + m^2$ can yield the same number for different pairs of (l, m) , e.g. $(7, 1)$, $(5, 5)$, $(1, 7)$ yield identical

$\omega_{lm} = 5\sqrt{2}\pi c/L$. For this example, then, there are degenerate 12 modes.

Expressing these modes as traveling waves gives, for example,

$$\begin{aligned} & 4 \cos(\kappa_l x_1) \cos(\kappa_m x_2) \cos(\omega_{lm} t) \\ = & \cos[\omega_{lm} t - (\kappa_l x_1 + \kappa_m x_2)] \\ & + \cos[\omega_{lm} t + (\kappa_l x_1 + \kappa_m x_2)] \\ & + \cos[\omega_{lm} t + (\kappa_l x_1 - \kappa_m x_2)] \\ & + \cos[\omega_{lm} t - (\kappa_l x_1 - \kappa_m x_2)]. \quad (27) \end{aligned}$$

As in Section 4.1, each of the degenerate traveling waves having equal amplitudes (on average) is equivalent to equipartition amongst the degenerate modes. These traveling waves having equal amplitudes means an equal amount of wave energy arriving from the specified directions. One could describe this as ‘discretely isotropic’, where ‘isotropic’ refers only to those discrete directions in which waves are allowed to travel. For the $\omega_{lm} = 5\sqrt{2}\pi c/L$ example given earlier, there are 12 of these directions and these directions are not evenly spaced in azimuth.

In the limit of short wavelength modes, $l^2 + m^2 \rightarrow \infty$, the number of degenerate traveling waves approaches ∞ and the directions become approximately evenly spaced. This fact can be understood with the following argument. Since $l^2 + m^2 = r^2$ is the equation of a circle in the (l, m) plane, solutions for modes can be described by the intersection of a circle of radius $r = \omega_{lm} L/(\pi c)$ with integer grid points. For example, $r = 5\sqrt{2}$ yields the 12 solutions $(\pm 7, \pm 1)$, $(\pm 5, \pm 5)$, $(\pm 1, \pm 7)$ discussed above, and the direction of incidence of those 12 waves are the 12 vectors given by the respective coordinates (see Fig. 1a). When $r \rightarrow \infty$, the circle has the opportunity to intersect many more grid points, and these grid points will become (approximately) evenly spaced in azimuth, especially if approximate intersection is allowed (i.e. approximately degenerate modes are counted, see Fig. 1b). Thus, we observe that in the limit of short wavelengths, equipartition of modes is equivalent to isotropic incidence of traveling waves.

One should note that if velocity structure is non-uniform (for example, if $c = c(\theta)$ has a dependence on azimuth, θ), then $r(\theta) = \omega_{lm} L/(\pi c(\theta))$ is no longer constant. For short wavelength modes, the intersection of this curve with grid points is no longer expected to be approximately evenly spaced in azimuth (see Fig. 1c). Equipartition amongst degenerate modes will therefore no longer imply isotropic incidence of waves. Instead, a very specific non-isotropic incidence of waves will imply equipartition, and only this special case will result in satisfying Eq. (23).

4.3 Waves on the Earth

Elastic waves on the Earth (including high frequency body waves) can be described in terms of (spheroidal and toroidal) normal modes (e.g. Dahlen & Tromp (1998)) so the formulation of Section 3 can still be used. As with the previous 2 examples, modes are discrete, with the longest period modes having a small number of (approximately) degenerate members (sometimes called multiplets). For these longest period modes (e.g. 400–4000 s), as in the examples of Section 4.2, equipartition of these degenerate modes would imply an ‘isotropic’ distribution of waves only in the discrete directions allowed by the Earth.

On the other hand, for short period modes (e.g. 10 s) whose wavelengths (e.g. 40 km) are very small compared to the Earth’s




Figure 1. Graphical depiction of solutions for degenerate (traveling wave) modes on a square. Each grid point corresponds to a mode. Intersections of the curve $r = \omega_{l,m} L / (\pi c)$ with grid points denote modes with the same degeneracy (denoted by stars). (a) When $r = 5\sqrt{2}$, the circle intersects 12 grid points. (b) When r is large, the modes are approximately evenly spaced. (c) When c is not constant, modes are no longer guaranteed to be evenly spaced. Intersections for cases (b) and (c) are approximate.

radius, the Earth can be approximated as flat and hence the situation is like that for the short wavelength modes on a periodic square of Section 4.2, except that there are multiple mode branches instead of just one. Thus, a large number of modes are degenerate and equipartition implies ‘nearly’ isotropic incidence of waves. As in Section 4.2, the qualifier ‘nearly’ is needed because the Earth has a (mostly weakly) heterogeneous velocity structure rather than being perfectly homogeneous. It should be noted that ‘nearly’ isotropic incidence does not necessarily imply equipartition if there are multiple mode branches that are (approximately) degenerate. For example, degenerate higher-order modes could have very low amplitudes and not be equipartitioned while degenerate fundamental modes could be equipartitioned, resulting in ‘nearly’ isotropic incidence but only partial equipartition. Time windowing as in Section 3.4 could be applied to consider only the fundamental branch.

Traditional ambient noise tomography applications on the Earth (e.g., Shapiro & Campillo 2004; Shapiro et al. 2005; Sabra et al. 2005) typically focus on relatively short-period waves (e.g. 5-40 s) excited by microseism energy. The implication of Section 3 is that if these short-period traveling waves are (nearly) isotropic, then these degenerate modes are equipartitioned and the cross correlation and Green’s function will be related by Eq. (23). If degenerate-mode waves are only partially equipartitioned (e.g. there exist higher order modes approximately degenerate with the fundamental branch of interest, as would be expected), a partial Green’s function may still be obtained from the cross correlation as described in Section 3.4 by use of time windowing, but quantifying the degree of success is difficult within the framework presented here. If waves are not equipartitioned, none of the analysis discussed here will have bearing on how cross correlation measurements can be related to the Green’s function. Analysis like that of Cox (1973), Harmon et al. (2008) or Tsai (2009), which describe a framework for examining non-isotropic incidence of waves with potentially non-uniform velocity structure, is better suited for quantifying the degree to which cross correlation measurements can still yield useful information related to the Green’s function. Since it is known that microseism energy is primarily generated in the oceans (which are not isotropically distributed), the success of recent noise tomography applications (e.g., Shapiro & Campillo 2004; Shapiro et al. 2005) is therefore still best understood through those alternative means. Nonetheless, the above analysis clarifies a number of points regarding noise cross correlation on the Earth, including the fact that equipartition need not be satisfied amongst all modes in order for a useful relationship between the cross correlation and Green’s function to exist.

5 CONCLUSIONS

In this work, we take a modal approach and demonstrate relationships between the cross correlation and the Green’s function, specifically examining the differences between the four cases where modes are equipartitioned or not and degenerate or not. We find that in certain cases waves need not be randomly excited and equipartition does not need to be strictly satisfied in order for a useful relationship to exist between the cross correlation and the Green’s function. Specifically, if no modes are degenerate then equipartition implies the standard relationship holds even with a completely deterministic system without random excitations; if

there are degenerate modes, equipartition must be accompanied by random phases for the standard relationship to hold; if modes are not equipartitioned but there is also no degeneracy, the relationship holds in a restricted sense to each frequency; finally, if modes are degenerate and not equipartitioned, partial Green's function phase information can sometimes be retrieved depending on whether partial equipartition exists. Applying this analysis to a few simple examples explains some aspects of noise cross correlation on the Earth, although quantitative analysis of the degree of success of noise tomography applications remains better understood through other means.

ACKNOWLEDGMENTS

The author thanks Y. Fan, L. Ramirez, S. Hartzell, T. Tada, D. E. McNamara and two anonymous reviewers for helpful comments. This research was supported by the Mendenhall Postdoctoral Fellowship program of the United States Geological Survey.

REFERENCES

- Cox, H., 1973. Spatial correlation in arbitrary noise fields with application to ambient sea noise, *J. Acoust. Soc. Am.*, **54**, 1289–1301.
- Dahlen, F. A. & Tromp, J., 1998. *Theoretical Global Seismology*, Princeton University Press, Princeton.
- Harmon, N., Gerstoft, P., Rychert, C. A., Abers, G. A., de la Cruz, M. S., & Fischer, K. M., 2008. Phase velocities from seismic noise using beamforming and cross correlation in Costa Rica and Nicaragua, *Geophys. Res. Lett.*, **35**.
- Lobkis, O. I. & Weaver, R. L., 2001. On the emergence of the Green's function in the correlations of a diffuse field, *J. Acoust. Soc. Am.*, **110**, 3011–3017.
- Roux, P., Sabra, K. G., Kuperman, W. A., & Roux, A., 2005. Ambient noise cross correlation in free space: Theoretical approach, *J. Acoust. Soc. Am.*, **117**, 79–84.
- Sabra, K. G., Gerstoft, P., Roux, P., Kuperman, W. A., & Fehler, M. C., 2005. Surface wave tomography from microseisms in Southern California, *Geophys. Res. Lett.*, **32**.
- Sanchez-Sesma, F. J. & Campillo, M., 2006. Retrieval of the green's function from cross correlation: The canonical elastic problem, *Bull. Seismol. Soc. Am.*, **96**, 1182–1191.
- Sanchez-Sesma, F. J., Perez-Ruiz, J. A., Campillo, M., & Luzon, F., 2006. Elastodynamic 2D Green function retrieval from cross-correlation: canonical inclusion problem, *Geophys. Res. Lett.*, **33**.
- Shapiro, N. M. & Campillo, M., 2004. Emergence of broadband Rayleigh waves from correlations of the ambient seismic noise, *Geophys. Res. Lett.*, **31**.
- Shapiro, N. M., Campillo, M., Stehly, L., & Ritzwoller, M. H., 2005. High-resolution surface-wave tomography from ambient seismic noise, *Science*, **307**, 1615–1618.
- Snieder, R., 2004. Extracting the Green's function from the correlation of coda waves: A derivation based on stationary phase, *Phys. Rev. E*, **69**, 1–8.
- Tsai, V. C., 2009. On establishing the accuracy of noise tomography travel-time measurements in a realistic medium, *Geophys. J. Int.*, **178**, 1555–1564.
- Wapenaar, K., 2004. Retrieving the elastodynamic Green's function of an arbitrary inhomogeneous medium by cross correlation, *Phys. Rev. Lett.*, **93**, 1–4.
- Wapenaar, K., Slob, E., & Snieder, R., 2006. Unified Green's function retrieval by cross correlation, *Phys. Rev. Lett.*, **97**.